

New and Old Approaches to Ice Sheet Modeling: Solid Earth Geophysics and the Cryosphere


**Jeremy N. Bassis
Douglas R. MacAyeal**

**Department of the Geophysical Sciences
University of Chicago
Scripps Institution of Oceanography,
University of California**

LANL Modeling Workshop: August 18-20

Solid earth geophysics and ice sheets


Mantle Convection



Geodynamics and ice sheets:

- Slow viscous flow
- Non-Newtonian fluid
- Temperature dependent viscosity
- Phase changes
- Brittle and ductile flow regimes

Solid earth geophysics and ice sheets



Geodynamics and ice sheets:

- Slow viscous flow
- Non-Newtonian fluid
- Temperature dependent viscosity
- Phase changes
- Brittle and ductile flow regimes


Low Peclet number:

$$Pe = \frac{\text{Mass Diffusion}}{\text{Mass Advection}}$$

Key differences:

- Ice sheets (nearly) barotropic
- Ice sheets don't conserve mass (mass added and removed)

Solid earth geophysics and ice sheets



Geodynamics and ice sheets:

- Slow viscous flow
- Non-Newtonian fluid
- Temperature dependent viscosity
- Phase changes
- Brittle and ductile flow regimes


Low Peclet number:

$$Pe = \frac{\text{Mass Diffusion}}{\text{Mass Advection}}$$


Key differences:

- Ice sheets (nearly) barotropic
- Ice sheets don't conserve mass (mass added and removed)


Ice sheet dynamics



Ice sheet dynamics



Ice sheet dynamics



Ice sheet dynamics

Shallow-ice approximation (SIA):

$$u(z) \sim 1 - \left(\frac{z_s - z}{h} \right)^{n+1}$$

ice-s

'Blatter'

$$u(z) \sim f(z)$$

ice-streams, outlet
glaciers, etc


Shallow-stream approximation (SSA):

$$u(z) \sim 1$$

ice-shelf

ocean

Bed rock/till



Ice sheet dynamics

Shallow-ice approximation (SIA):

$$u(z) \sim 1 - \left(\frac{z_s - z}{h} \right)^{n+1}$$

ice-s

Goal: Describe transition from SIA to SSA

'Blatter'

$$u(z) \sim f(z)$$

Shallow-stream approximation (SSA):


$$u(z) \sim 1$$

ice-streams, outlet
glaciers, etc

ice-shelf

ocean

Bed rock/till




A simple approximation

Multiple (equivalent) approaches to mechanics:
(Newtonian, Lagrangian, Poisson Brackets, . . .)

Start with Hamilton's Principle:

$$L = \int_{t_1}^{t_2} (\text{kinetic energy} - \text{potential energy}) dt$$

$$\delta L = 0$$




A simple approximation

Multiple (equivalent) approaches to mechanics:
(Newtonian, Lagrangian, Poisson Brackets, . . .)

Start with Hamilton's Principle:

$$L = \int_{t_1}^{t_2} (\text{kinetic energy} - \text{potential energy}) dt$$

$$\delta L = 0$$



A simple approximation

Multiple (equivalent) approaches to mechanics:
(Newtonian, Lagrangian, Poisson Brackets, . . .)


Start with Hamilton's Principle:

$$L = \int_{t_1}^{t_2} (\text{kinetic energy} - \text{potential energy}) dt$$

$$\delta L = 0$$

$$dU = \text{constant entropy} + dW$$

$$dW = PdV + d\Gamma_{\text{visc}} + d\Gamma_{\text{fric}}$$



A simple approximation

Multiple (equivalent) approaches to mechanics:
(Newtonian, Lagrangian, Poisson Brackets, . . .)

Start with Hamilton's Principle:

$$L = \int_{t_1}^{t_2} (\text{kinetic energy} - \text{potential energy}) dt$$

$$\delta L = 0$$

$$dU = \cancel{TdS} + dW$$

constant entropy

$$dW = PdV + d\Gamma_{\text{visc}} + d\Gamma_{\text{fric}}$$

Dissipation of gravitational potential = viscous + frictional dissipation energy

$L[u]$ = functional of u to be minimized

Bed rock/till

ice-sheet

ice-shelf

ocean

A simple approximation: Rayleigh-Ritz Method


Assume velocity of the form:

$$u(x,z) = a_0 + a_1\phi_1 + a_2\phi_2 + \dots$$

Require:

$$u(x,z) = \text{SIA}$$

$$u(x,z) = \text{SSA}$$



A simple approximation: Rayleigh-Ritz Method

Assume velocity of the form:

$$u(x,z) = a_0 + a_1\phi_1 + a_2\phi_2 + \dots$$

Require:

$$u(x,z) = \text{SIA}$$

$$u(x,z) = \text{SSA}$$

Two term expansion:

$$u(x,z) = a_0(x) + a_1(x) \left[1 - \left(\frac{z_s - z}{h} \right)^{n+1} \right]$$

Satisfy Basal BC:


$$u(x,z) = U(x) \left\{ \frac{\varepsilon(x)}{n+1} \left[1 - \left(\frac{z_s - z}{h} \right)^{n+1} \right] + 1 \right\}$$

ice-sheet

ocean

Bed rock/till

A simple approximation: Rayleigh-Ritz Method



(Non-dimensional) Diagnostic and Prognostic Equations


$$\frac{\partial}{\partial x} \left[c_1(x) \frac{\partial U}{\partial x} \right] + c_2(x)U = h \frac{\partial z_s}{\partial x} + \epsilon \delta^{-2} \beta' U$$

Transition from SIA to SSA introduces
'quasi-boundary layers'

$$\frac{\partial h}{\partial t} + \frac{\partial}{\partial x} [h (u - u_{SIA})] - \frac{\partial}{\partial x} \left(D \frac{\partial h}{\partial x} \right) = M$$

Non-shallow ice flux shallow ice flux

Time Integration: Mixed Eulerian-Lagrangian Methods




Time Integration: Mixed Eulerian-Lagrangian Methods

Allow integration points to advect as Lagrangian tracers

Create/destroy tracers to preserve accuracy

Can view tracers as discrete mass - **mass is conserved exactly**




Time Integration: Mixed Eulerian-Lagrangian Methods

Allow integration points to advect as Lagrangian tracers

Create/destroy tracers to preserve accuracy

Can view tracers as discrete mass - **mass is conserved exactly**




Time Integration: Mixed Eulerian-Lagrangian Methods

Allow integration points to advect as Lagrangian tracers

Create/destroy tracers to preserve accuracy

Can view tracers as discrete mass - **mass is conserved exactly**



nodes migrate

Advantages:

Lagrangian advection of information (no fake diffusion)

Re-meshing is interpolation free (nodes don't get moved)

Easy to deal with evolving free surfaces

Disadvantages:

Resolution migrates


Ad-hoc rules to add/subtract nodal points

Boundary conditions can be problematic

Need about 2x as many grid nodes

Grounding line migration: 'Particle Method'

- Represent grounding line as a quasi-particle - a 'groundon'
- Grounding line migration - find groundon trajectory




- Evolution equation for each groundon:

$$\frac{dg_1}{dt} = f_1(t)$$

- Solve simultaneously with advection-diffusion eqn.

Grounding line migration: 'Enthalpy Method'

- Analogy with diffusion problems with moving phase boundaries
- Consider floating ice and grounded ice to be different phases



Boundary has to advance/retreat one discrete grid point at a time

Introduce parameter f (varies from 0 to 1) = $\frac{\text{Volume of ice in grid}}{\text{Volume to ground entire cell}}$


Track flux of ice into each grid over time - removes some of hysteresis problems

Which ice dynamics approximation is best?

- Depends on **question** asked **and** **data** available
- A model is an improvement if it explains more data
- But . . . also need to account for increased number of degrees of freedom

Which ice dynamics approximation is best?


- Depends on **question** asked **and** **data** available
- A model is an improvement if it explains more data
- But . . . also need to account for increased number of degrees of freedom



- Need to consider model improvement **per degree of freedom**

Which ice dynamics approximation is best?


- Depends on **question** asked **and** **data** available
- A model is an improvement if it explains more data
- But . . . also need to account for increased number of degrees of freedom



- Need to consider model improvement **per degree of freedom**
- How much of observation-space is explained by different approximations

Which ice dynamics approximation is best?


- Depends on **question** asked **and** **data** available
- A model is an improvement if it explains more data
- But . . . also need to account for increased number of degrees of freedom



- Need to consider model improvement **per degree of freedom**
- How much of observation-space is explained by different approximations

Which ice dynamics approximation is best?


- Depends on **question** asked **and** **data** available
- A model is an improvement if it explains more data
- But . . . also need to account for increased number of degrees of freedom



- Need to consider model improvement **per degree of freedom**
- How much of observation-space is explained by different approximations

Which ice dynamics approximation is best?


- Depends on **question** asked **and** **data** available
- A model is an improvement if it explains more data
- But . . . also need to account for increased number of degrees of freedom



- Need to consider model improvement **per degree of freedom**
- How much of observation-space is explained by different approximations

Which ice dynamics approximation is best?



- Depends on **question** asked **and** **data** available
- A model is an improvement if it explains more data
- But . . . also need to account for increased number of degrees of freedom



- Need to consider model improvement **per degree of freedom**
- How much of observation-space is explained by different approximations


An Example: Steady-state ice sheet profiles

$$\beta = 10^{14} \text{ Pa}\cdot\text{s}\cdot\text{m}^{-1}$$



An Example: Steady-state ice sheet profiles


$$\beta = 10^{14} \text{ Pa} \cdot \text{s} \cdot \text{m}^{-1}$$



An Example: Steady-state ice sheet profiles

$$\beta = 10^{14} \text{ Pa} \cdot \text{s} \cdot \text{m}^{-1}$$


near terminus strain rate




An Example: Steady-state ice sheet profiles

$$\beta = 10^{14} \text{ Pa} \cdot \text{s} \cdot \text{m}^{-1}$$


near terminus strain rate



An Example: Steady-state ice sheet profiles




An Example: Steady-state ice sheet profiles




An Example: Steady-state ice sheet profiles

'ice-stream'




An Example: Steady-state ice sheet profiles


'ice-stream'




Steady-state ice sheet profiles




Steady-state ice sheet profiles



Steady-state ice sheet profiles




Steady-state ice sheet profiles




Back to explanation space . . .

- Depends on **question** asked **and** **data** available
- A model is an improvement if it explains more data
- But . . . also need to account for increased number of degrees of freedom




Back to explanation space . . .

- Depends on **question** asked **and** **data** available
- A model is an improvement if it explains more data
- But . . . also need to account for increased number of degrees of freedom



Back to explanation space . . .

- Depends on **question** asked **and** **data** available
- A model is an improvement if it explains more data
- But . . . also need to account for increased number of degrees of freedom



Challenge: Identify and parameterize the 'barrier' processes that limit our understanding

Deux ex machina?

Already developed
and in use

‘SIA/SSA’

Coarse grids, integration over hundreds
of thousands of years

Heavy development
now

‘Blatter’

Moderate grids, century to thousand
year time-scale integration

Highest development
cost

‘Full Stokes’

High resolution grids, decade to cen-
tury time-scale integration

Deux ex machina?

Already developed
and in use

`SIA/SSA'

Coarse grids, integration over hundreds
of thousands of years

Heavy development
now

`Blatter'

Moderate grids, century to thousand
year time-scale integration

Highest development
cost

`Full Stokes'

High resolution grids, decade to cen-
tury time-scale integration

